

COUNTRY FIELD FORAGE 2487 ORGANIC

Clover grass annual



Profile

The non winterhardy mixture is ideal for several cuts and delivers good yields. COUNTRY Field Forage 2487 is optimal for spring sowing. The combination of different types of annual clover and ryegrass provide a protein rich quality forage and ensure the supply of nutrients.

- ✓ With annual clovers
- ✓ Leafy growth
- ✓ Flexible in use



Components

30%	Italian ryegrass Dorike
30%	Lolium westerwoldicum Arnoldo
20%	Egyptian clover Bluegold
20%	Persian clover Resal

* also suitable for intercropping, yield in 1st cut above average

Sowing pattern

New sowing

Locations

Dry	<div><div>+</div><div></div><div></div></div>
Normal	<div><div>+</div><div>+</div><div>+</div></div>
Wet	<div><div>+</div><div>+</div><div>+</div></div>
Peat Soil	<div><div>+</div><div></div><div></div></div>
Altitude	<div><div>+</div><div>+</div><div></div></div>

Usage

Grazing	<div><div>+</div><div>+</div><div></div></div>
Grazing and cutting	<div><div>+</div><div>+</div><div></div></div>
Cutting	<div><div>+</div><div>+</div><div>+</div></div>
Extensive usage	<div><div>+</div><div>+</div><div></div></div>

Fodder values

Energy density	<div><div>+</div><div>+</div><div>+</div></div>
Digestibility	<div><div>+</div><div>+</div><div>+</div></div>
Protein	<div><div>+</div><div>+</div><div>+</div></div>

+++ = highly suitable | ++ = suitable | + = conditionally suitable | - = not given

Cultivation notes

New sowing	35 - 40 kg/ha
Sowing time	from April to August
Usage per year	1 - 5

All organic mixtures are produced with 100 % organic content, if possible, in accordance with the EU Organic Farming Regulation 2018/848. For mixtures with at least 70% organic content, the farmer must obtain approval from his inspection body for the use of the conventional content before sowing, starting with production in January 2022. We aim to use at least 2 varieties per ripening group for mix proportions of more than 20 % of one species and for German ryegrass. The choice of varieties may vary depending on production. If individual species or varieties are not available, they are replaced by equivalent ones. The seed strengths are calculated for an optimal seedbed; in case of poor external conditions - such as insufficient seedbed, late sowing date, insufficient nutrient supply and unfavourable weather - the seed strengths should be adjusted upwards.

All specified informations, recommendations and representations mentioned here are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG excludes adhesion for damage or claims for damages, resulting of the use for the variety specified in this description. Mixture compositions may change if individual varieties are not available. As of 07/2023. Subject to change without notice.