

COUNTRY GRASSLAND 2463 ORGANIC

Overseeding dry sites



Profile

Due to the balanced composition of species and their maturity groups, this mixture ensures optimal utilization of winter moisture and thus puts the focus on the 1st cut. The use of Festulolium increases the drought tolerance of the mixture, resulting in improved yield potential. Perennial ryegrass and Festulolium are ideal for reseeding due to rapid youth development and competitive vigor.

- ✓ Drought-tolerant Festulolium
- ✓ Exploitation of winter moisture
- ✓ Emphasis on first cut



Components

30%	Lolium perenne (early) Karatos, Mirtello
23%	Lolium perenne Explosion
20%	Lolium perenne Valerio
20%	Festulolium Fedoro
7%	Trifolium repens Abercrest, Liflex

Sowing pattern

New sowing, Overseeding

Locations

Dry	<div><div>+</div><div>+</div><div>+</div></div>
Normal	<div><div>+</div><div>+</div><div></div></div>
Wet	<div><div>+</div><div></div><div></div></div>
Peat Soil	<div><div>+</div><div></div><div></div></div>
Altitude	<div><div>+</div><div>+</div><div></div></div>

Usage

Grazing	<div><div>+</div><div>+</div><div></div></div>
Grazing and cutting	<div><div>+</div><div>+</div><div>+</div></div>
Cutting	<div><div>+</div><div>+</div><div>+</div></div>
Extensive usage	<div><div>+</div><div></div><div></div></div>

+++ = highly suitable | ++ = suitable | + = conditionally suitable | - = not given

Cultivation notes

New sowing	35 - 40 kg/ha
Overseeding	20 - 25 kg/ha
Overseeding, several times per year	7 - 10 kg/ha
Sowing time	from March to September
Usage per year	3 - 6

All organic mixtures are produced with 100 % organic content, if possible, in accordance with the EU Organic Farming Regulation 2018/848. For mixtures with at least 70% organic content, the farmer must obtain approval from his inspection body for the use of the conventional content before sowing, starting with production in January 2022. We aim to use at least 2 varieties per ripening group for mix proportions of more than 20 % of one species and for German ryegrass. The choice of varieties may vary depending on production. If individual species or varieties are not available, they are replaced by equivalent ones. The seed strengths are calculated for an optimal seedbed; in case of poor external conditions - such as insufficient seedbed, late sowing date, insufficient nutrient supply and unfavourable weather - the seed strengths should be adjusted upwards.

All specified informations, recommendations and representations mentioned here are made to the best of our knowledge and belief, but without guarantee of completeness or accuracy. We cannot guarantee that the properties described are repeatable. All information is provided as an aid to decision-making. Deutsche Saatveredelung AG excludes adhesion for damage or claims for damages, resulting of the use for the variety specified in this description. Mixture compositions may change if individual varieties are not available. As of 07/2023. Subject to change without notice.